animal

Effects of dietary calcium and phosphorus deficiency and subsequent recovery on broiler chicken growth performance and bone characteristics

A. S. Valable^{1,2}, A. Narcy², M. J. Duclos², C. Pomar³, G. Page⁴, Z. Nasir⁴, M. Magnin⁵ and M. P. Létourneau-Montminy^{1†}

¹Animal Sciences Department, Université Laval, 2425 rue de l'Agriculture, Québec, QC, Canada G1V 0A6; ²URA, INRA, 37380, Nouzilly, France; ³Agriculture and Agri-Food Canada, Dairy and Swine Research and Development Center, Sherbrooke, QC, Canada J1M 123; ⁴Trouw Nutrition Agresearch, 150 Research Lane, Guelph, ON, Canada, N1G 472; ⁵MiXscience, Centre d'affaires Odyssée, ZAC Cicé Blossac, 35172 BRUZ, France

(Received 19 April 2017; Accepted 5 September 2017; First published online 4 December 2017)

The ability of birds to modify dietary phosphorus utilisation when fed with low-phosphorus and calcium (Ca) diets was studied using different sequences of dietary phosphorus and Ca restriction (depletion) and recovery (repletion) during the grower and the finisher phases. A total of 3600 Ross 708 broilers were randomly divided into 10 replicate pens per treatment (60 per pen, six pens per block). Chicks were fed a common starter diet from days 0 to 10, then a grower control diet (C: 0.90% Ca, 0.39% non-phytate phosphorus, nPP), mid-level diet (M: 0.71% Ca, 0.35% nPP) or low Ca and nPP diet (L: 0.60% Ca, 0.30% nPP) from days 11 to 21, followed by a finisher diet C, M or L containing, respectively, 0.85%, 0.57% or 0.48% Ca and 0.35%, 0.29% or 0.24% nPP from days 22 to 37. Six treatment sequences were tested: CC, MM, LL, ML, LC and LM. Bone mineral content by dualenergy X-ray, tibia ash, toe ash weight and tibia breaking strength were measured on days 21 and 37. No significant effect was observed on growth performance throughout the experiment. Diet L reduced bone mineral content, breaking strength, tibia and toe ash by 9%, 13%, 11% and 10%, respectively, on day 21 (compared with diet C, for linear effect, P < 0.05). On day 37, bone mineral content, breaking strength, tibia and toe ash remained lower compared with control values (CC v. MM v. LL, P < 0.05 for linear and quadratic effects). Mineral depletion duration (ML v. LL) did not affect bone mineral status. Replenishing with the C diet during the finisher phase (LC) restored bone mineral content, tibia ash and toe ash weight better than the M diet did, but not to control levels (CC v. LC v. LM, for linear effect, P < 0.05). These results confirm that dietary Ca and nPP may be reduced in the grower phase without affecting final growth performance or breaking strength as long as the finisher diet contains sufficient *Ca* and *nPP*. The practical applications of this strategy require further study in order to optimise the depletion and repletion steps.

Keywords: broiler chickens, feeding strategy, calcium, phosphorus, adaptation

Implications

A broiler chicken grower diet containing less calcium and phosphorus than recommended (depletion) affects bone mineral status but not growth performance. Bone mineralisation is restored at least partially when the finisher diet contains adequate amounts of calcium and phosphorus (repletion). Calcium and phosphorus utilisation efficiency appears to increase under depletion conditions. This suggests that phosphorus ingestion and excretion can be reduced and that the sustainability of broiler production can thus be improved.

Introduction

Phosphorus utilisation is of increasing concern regarding the sustainability of broilers' production, considering high phosphorus excretion can lead to eutrophication. Phosphorus represents an important nutrient constraining feed costs. Several nutritional strategies have been developed to improve its utilisation. Yan *et al.* (2005) demonstrated that chickens fed diets deficient in calcium (Ca) and non-phytate phosphorus (nPP) during an early phase of growth (depletion) and later fed diet containing adequate levels of these minerals (repletion) exhibited better phosphorus and Ca utilisation and bone mineralisation. In fact, birds seem to respond to a decrease in Ca and nPP dietary

[†] E-mail: marie-pierre.letourneau-montminy@fsaa.ulaval.ca

content by increasing expression of mRNA encoding Ca transporters (Centeno et al., 2004; Bar, 2009) and phosphorus transporters (Ashwell and Angel, 2010) in the small intestine. Some authors have suggested that recommendations made by National Research Council (NRC) (1994) regarding dietary Ca and phosphorus for broiler chickens might exceed actual requirements (Angel et al., 2006; Létourneau-Montminy et al., 2010). In a meta-analysis, Létourneau-Montminy et al. (2010) have shown that feeding 21-day-old broilers 6.0 g of Ca and 3.0 g of nPP/kg of BW did not affect growth performance. Concomitant reduction of dietary nPP and Ca maintains phosphorus availability in the digestive tract. However, this can lead to poor bone mineralisation and thus impairing animal welfare or increased processing losses (Applegate and Angel, 2008). We focused on the grower phase for the initial depletion as a reduction of phosphorus and Ca intake during an early phase could lead to imprinting allowing birds to be more efficient for the entire period. More precisely, we expected that the initial depletion could allow a reduction in phosphorus intake later during the finisher phase. The results would allow the highest economic and environmental gain as birds consume two-thirds of feed during the finisher phase. We hypothesised that: (1) we could lower the nPP and Ca levels of the diet compared with NRC (1994) recommendations without impairing animal performance, (2) the depletion will reduce bone mineralisation and induce regulations which in turn will increase phosphorus and Ca utilisation efficiency, (3) broilers previously depleted will deposite more Ca and phosphorus (g/d) in the bone allowing them to catch-up in bone mineralisation at the control levels during repletion. Thus, this study was designed to test the effect of several Ca and nPP depletion degree and duration in order to improve Ca and phosphorus utilisation, and the levels of dietary Ca and nPP required subsequently in order to replenish the bone mineral deficit thus created.

Material and methods

Experimental diets

Except for Ca and phosphorus, all nutritional requirements of broiler chickens (NRC, 1994) were met or exceeded by the maize, soya bean meal-based diets formulated (Table 1). During the starter period (0 to 10 days of age), the chicks received a diet meeting all nutritional requirements (NRC, 1994). During the grower period (11 to 21 days of age), they received the control diet (C, 0.90% Ca and 0.39% nPP), the mid-level diet (M, 0.71% Ca and 0.35% nPP) or the low-level diet (L, 0.60% Ca and 0.30% nPP). During the finisher phase (22 to 37 days of age), Ca and nPP contents were, respectively, 0.85%, 0.57% and 0.48% and 0.35%, 0.29% and 0.24% in the C. M and L diets. The C diet contained Ca and nPP at the levels used in standard commercial conditions in Canada, whereas the L diet was based on previous studies (Yan et al., 2005; Létourneau-Montminy et al., 2010; Rousseau et al., 2016). The M diet was intermediate between the C and L diet and had the same Ca:nPP ratio as the L diet. Grower and finisher diets were combined to form six dietary

treatment sequences: CC, MM, LC, LM, ML and LL. The first treatment (CC) is the North American positive control whereas the second is the French positive control (MM) according to Létourneau-Montminy *et al.* (2010) and Rousseau *et al.* (2013 and 2016). The third (LC) and fourth (LM) dietary sequences have been built to test depletion and repletion strategies. When comparing them it will be possible to set the level (C or M) allowing a catch-up in bone mineralisation to reach the level of the two first dietary sequences. The fifth sequence (ML) will informed about the dietary Ca and phosphorus reduction that is possible to perform during finishing phase (22 to 35 days) for which few information is available in literature. Finally, the last dietary treatment represents the negative control (LL).

On each phase, one basal diet was manufactured and then divided in three mix corresponding to the different diets. In each mix, Ca and phosphorus sources (calcium carbonate and di-calcium phosphate), maize (19.6 and 21.7 g/kg during grower and finisher phase) and bentonite were added to obtain the different Ca and nPP levels required. Diets were fed as medium crumb during starter phase and as short pellets during grower and finisher phases. Birds had free access to feed and water throughout the experiment.

Birds and management

Animals were cared for according to the guidelines of the Canadian Council on Animal Care (2009). A total of 3600 1-day-old male Ross 708 broiler chicks were used. On arrival, chicks were assigned randomly to pens, 60 chicks/pen. The study fit a random block design consisting of 10 blocks of six pens each. The room temperature was settled at 32°C upon arrival and decreased progressively to reach 24°C at day 35 with a 23L:1D schedule from 0 to 7 day-of-age and 20L:1D schedule after day 7. Total chick weight in the pen and mean chick body mass were determined at 0, 10, 21 and 37 days of age. Feed intake per pen was measured for each period. Morbidity and mortality were recorded. Barn temperature and relative humidity were recorded daily. At the end of each period, one or two birds per pen were euthanised using CO₂ to sampled 10 birds/treatment. They were scanned dorsally in prone position by dual-energy X-ray absorptiometry (DXA) according to the method used by Schreiweis et al. (2005) to determine total bone mineral content and total body lean and fat contents. Schreiweis et al. (2005) had validated the use of DEXA to assess bone integrity in birds compared with tibia ash and breaking strength. The acquisitions were realised using the small animal mode as recommended by Mitchell et al. (1997) (DPX-L; LUNAR Prodigy Advance Corp., Madison, WI, USA) in a study comparing DXA results with carcass dissection measurements for bone mineral content and total body lean and fat contents. Tibias and toes were then collected for mineral analysis.

Laboratory analysis

Representative samples of the diets were taken upon delivery and once weekly. Samples of each feed were mixed together weekly and at the end of the experiment to obtain a

 Table 1 Composition of experimental diets fed to broiler chickens aged 11 to 37 days

		Grower phase diet	1	Finisher phase diet ¹				
Diets	C	М	L	С	М	L		
Ingredient (g/1050 kg) ²								
Maize	610.2	610.2	610.2	622.8	622.8	622.8		
Wheat	_	_	_	52.6	52.6	52.6		
Maize DDGS ³	9.7	9.7	9.7	-	-	-		
Soya bean meal	259.9	259.9	259.9	200.0	200.0	200.0		
Feather meal	31.6	31.6	31.6	31.6	31.6	31.6		
Fat	66.0	66.0	66.0	57.9	57.9	57.9		
Maize gluten meal	26.3	26.3	26.3	41.8	41.8	41.8		
Calcium carbonate	15.7	11.7	9.7	15.4	9.4	7.9		
Di-calcium phosphate	13.2	11.2	8.6	11.4	8.5	6.1		
Bentonite	_	6.0	10.6	-	8.9	12.8		
Sodium carbonate	3.0	3.0	3.0	3.5	3.5	3.5		
Choline chloride (70%)	0.7	0.7	0.7	0.9	0.9	0.9		
Salt	1.9	1.9	1.9	1.4	1.4	1.4		
Premix ⁴	4.6	4.6	4.6	4.6	4.6	4.6		
L-Lysine	4.0	4.0	4.0	3.8	3.8	3.8		
DL-Methionine	2.4	2.4	2.4	1.8	1.8	1.8		
Threonine	0.8	0.8	0.8	0.5	0.5	0.5		
Calculated composition ⁵								
Dry matter	87.1	87.0	87.0	87.1	87.0	87.0		
Metabolisable energy (MJ/kg)	12.5	12.5	12.5	12.7	12.7	12.7		
Crude fat	8.6	8.6	8.6	7.8	7.8	7.8		
СР	20.2	20.2	20.2	18.9	18.9	18.9		
Crude fibre	1.9	1.9	1.9	1.8	1.8	1.8		
Calcium (Ca)	0.90 (0.95)	0.71 (0.75)	0.60 (0.65)	0.85 (0.85)	0.58 (0.57)	0.48 (0.54)		
Total phosphorus	0.60 (0.62)	0.56 (0.58)	0.51 (0.53)	0.54 (0.53)	0.48 (0.49)	0.43 (0.47)		
Non-phytate phosphorus (nPP)	0.40	0.35	0.30	0.35	0.29	0.24		
Ca:nPP ratio	2.25	2.00	2.00	2.43	2.00	2.00		

C = standard commercial levels of Ca and nPP (positive control); M = mid-levels of Ca and nPP; L = low levels.

¹Days 11 to 21 and days 22 to 37.

²1000 kg of masterbatch diet, 50 kg to adjust Ca and nPP levels.

³Maize distillers dried grains.

⁴Per kg: 6487 IU retinol; 2366 IU cholecalciferol; 20.8 IU $D_{L-\alpha}$ -tocopherol; 3 mg menadione; 2 mg thiamine; 6 mg riboflavin; 78 mg niacin; 14 mg pantothenate; 3 mg pyridoxine; 104 mcg biotin; 1 mg folic acid; 14 mcg cyanocobalamin; 35 mg Fe; 65 mg Cu; 92 mg Mn; 92 mg Zn; 2 mg I. ⁵Calculated from NRC (1994). All values except metabolisable energy are percentages. Analysed values in brackets.

representative composite sample and analysed using the Association of Analytical Communities official methods for dry matter, protein and ash (Methods 930.15, 990.03 and 942.05). Ash was then solubilized using HCl in order to determine dietary Ca and phosphorus by inductively coupled plasma mass spectometry (Method 985.01). The right and left tibias were freed from adhering tissue. The middle left toe was separated between the second and the third metatarsal bone without removing either nail or skin, then cleaned. The left tibia and toes were weighed, dried at 110°C for 24 h and then placed in a muffle furnace (Lindberg/Blue M Vacuum oven; Thermo Scientific, Waltham, MA, USA) at a temperature ramped from 200°C to 600°C over 6 h followed by 48 h at 600°C. The right tibia was used to measure breaking strength (Chatillon TCM 201; Wagner Instrument, Greenwich, CT, USA).

Calculation and statistical analysis

Bone mineral content gain was calculated by dietary treatments for each bird as the bone mineral content at the end of the phase minus the mean for the same dietary treatment at the beginning of the phase. The Ca and nPP relative transfer to whole-body, tibia and toe were calculated separately for the bone criteria: bone mineral content gain, tibia and toe ash weight. It was defined as the ratio between the value of each bone criterion (g) and the Ca or nPP ingested (g) during the corresponding period. Ingested Ca and nPP were calculated from the average daily feed intake over the period for the corresponding pen. The pen was the experimental unit for growth performance and the bird was the experimental unit for bone mineralisation. Dietary sequences were included in the model as a fixed effect and the block as a random effect. ANOVA were performed using the MIXED procedure of SAS (Statistical Analysis System Institute Inc., 2003) after the normality of the variables had been checked using the Shapiro–Wilk test. Orthogonal contrasts with linear (Lin) and quadratic (Qua) effects were used in the grower phase (C v. M v. L) and for the overall experiment to study the extent of the depletion in dietary nPP and Ca levels (CC v. MM v. LL),

Valable, Narcy, Duclos, Pomar, Page, Nasir, Magnin and Létourneau-Montminy

the impact of the depletion duration (ML v. LL), and the effect of the levels of dietary nPP and Ca during the finisher phase (CC v. LC v. LM). Linear regression between tibia, toe and whole-body criteria was conducted within each phase. Differences were considered significant at P < 0.05.

Results

Analysed phosphorus and Ca concentrations were in accordance with expected values.

Grower phase

Diet had no effect on growth performance measured on day 21 (Table 2). Birds fed the L diet had a higher body fat content (g/100 g BW) than those fed the C diet (+18%, Lin, P = 0.001), whereas lean content (g/100 g BW) decreased with a decreased of dietary Ca and nPP (-3%, Lin, P < 0.001). A reduction of 33% of Ca and of 23% of nPP content reduced bone mineral content and gain (-9% and -11%, Lin, P < 0.001), breaking strength (-13%, Lin, P = 0.01), tibia and toe ash weight (-11% and -10%, Lin, P < 0.001). Lowering the level of Ca and nPP in the diets improved the Ca relative transfer to tibia (Lin, P < 0.001) and

nPP relative transfer to whole-body, tibia and toe (Lin, P < 0.001). Based on bone mineral content and toe ash weight, C diet lead to lower Ca relative transfer to whole-body and toe compared with M and L diets (Lin, P < 0.001, Qua, P < 0.05).

Finisher phase

Impact of reducing dietary phosphorus and calcium. The effect of the reduced intake of Ca and phosphorus is shown by the contrast CC v. MM v. LL. Growth performance were unaffected (Table 3). The LL treatment resulted in a lower bone mineral content gain (Lin, P < 0.001). The MM and LL treatments both led to lower bone mineral content, tibia breaking strength, tibia ash and toe ash weights than did the control treatment (Lin, Qua, P < 0.01) but higher Ca relative transfer to whole-body, tibia and toe (Lin, Qua, P < 0.001) and higher nPP relative transfer to toe (Lin, P = 0.01, Qua, P < 0.05). The duration of the depleting treatment (ML v. LL contrast) had no effect on performance.

Impact of dietary phosphorus and calcium during the repletion period. The effect of replenishing the two minerals is shown by the contrast CC v. LC v. LM. Growth performance were not affected (Table 3). On day 37, breaking strength

Table 2 Effect of calcium (Ca) and non-phytate phosphorus (nPP) on broiler chicken growth performance and bone mineralisation during the grower phase (days 11 to 21)

					P value ¹	
		Diet ²		C <i>v</i> . M <i>v</i> . L		
Response criteria	С	М	L	SEM	Lin	Qua
Performance						
Final BW (g)	920.6	922.5	920.0	5.9	0.93	0.68
ADG (g/d)	61.79	61.94	61.72	0.44	0.89	0.66
ADFI (g/d)	84.45	84.02	84.45	0.58	0.86	0.43
FCR	1.387	1.379	1.390	0.013	0.75	0.15
Body fat (g/100 g BW)	11.34	12.65	13.39	0.56	0.001	0.39
Body lean mass (g/100 g BW)	87.16	85.63	85.00	0.58	<0.001	0.37
Mortality (%)	0.33	0.75	0.89	0.36	0.18	0.67
Bone mineralisation						
Bone mineral content (g)	15.68	15.20	14.24	0.29	<0.001	0.51
Bone mineral content gain (g/d)	13.10	12.61	11.66	0.29	<0.001	0.51
Tibia breaking strength (N)	195.6	174.5	171.2	6.8	0.01	0.28
Tibia ash weight (g)	1.03	0.98	0.92	0.02	<0.001	0.99
Toe ash weight (g)	0.131	0.125	0.118	0.003	<0.001	0.78
Ca relative transfer						
Bone mineral content gain (g/g)	0.77	0.94	0.99	0.02	<0.001	0.03
Tibia ash weight (g/g)	0.061	0.074	0.080	0.002	<0.001	0.08
Toe ash weight (g/g)	0.0079	0.0096	0.0103	0.0002	<0.001	0.03
nPP relative transfer						
Bone mineral content gain (g/g)	1.74	1.85	1.96	0.05	0.001	0.91
Tibia ash weight (g/g)	0.139	0.146	0.158	0.004	<0.001	0.48
Toe ash weight (g/g)	0.0178	0.0189	0.0204	0.0005	<0.001	0.62

C = standard commercial levels of Ca and nPP (positive control); M = mid-levels of Ca and nPP; L = low levels; Lin = linear effect; Qua = quadratic effect; ADG = average daily gain; ADFI = average daily feed intake; FCR = feed conversion ratio.

For growth performance, n = 10 pens for C, n = 20 for M, n = 30 for L. For bone mineralisation, n = 10 birds/treatment.

²See Table 1.

Table 3 Effect of calcium (Ca) and non-phytate phosphorus (nPP) on broiler chicken growth performance and bone mineralisation during the finishing phase (days 22 to 37)

								P value ¹				
									Depletior	Reple	Repletion	
			Di	et ²				CC v. MM v. LL ML v. LL CC v. LC			C <i>v</i> . LM	
Parameters	СС	ММ	LL	ML	LC	LM	SEM	Lin	Qua	Lin	Lin	Qua
Performance												
Final BW (g)	2468	2450	2441	2444	2463	2483	30	0.43	0.57	0.86	0.61	0.64
ADG (g/d)	95.6	93.7	94.2	94.6	95.2	97.0	1.6	0.35	0.73	0.67	0.38	0.41
ADFI (g/d)	163.4	166.0	163.5	165.5	163.8	165.9	2.5	0.96	0.56	0.83	0.32	0.71
FCR	1.708	1.774	1.736	1.755	1.722	1.710	0.025	0.26	0.81	0.45	0.95	0.57
Body fat (g/100 g BW)	10.32	10.88	10.80	10.96	10.89	10.51	0.36	0.69	0.26	0.52	0.25	0.40
Body lean mass (g/100 g BW)	85.81	85.35	85.35	85.34	85.75	85.70	0.37	0.82	0.34	0.45	0.91	0.32
Mortality (%)	0.93	0.95	0.86	0.75	0.75	0.94	0.58	0.90	0.86	0.73	0.98	0.71
Bone mineralisation												
Bone mineral content (g)	37.3	34.9	31.8	33.2	36.6	33.9	1.2	< 0.001	< 0.001	0.12	0.01	0.30
Bone mineral content gain (g/d)	21.81	19.99	16.93	17.69	22.51	19.57	0.07	<0.001	0.09	0.48	0.08	0.10
Tibia breaking strength (N)	345	311	281	331	341	314	21	0.002	0.01	0.30	0.14	0.54
Tibia ash weight (g)	2.81	2.54	2.31	2.50	2.67	2.48	0.09	< 0.001	< 0.001	0.63	0.001	0.74
Toe ash weight (g)	0.318	0.301	0.283	0.289	0.323	0.298	0.007	< 0.001	0.002	0.22	0.05	0.08
Ca relative transfer												
Bone mineral content gain	0.98	1.30	1.24	1.26	1.04	1.30	0.04	< 0.001	< 0.001	0.65	< 0.0001	0.02
(g/g)												
Tibia ash weight (g/g)	0.127	0.168	0.163	0.174	0.120	0.164	0.006	< 0.001	0.001	0.22	<0.001	< 0.001
Toe ash weight (g/g)	0.014	0.020	0.020	0.020	0.015	0.020	0.001	< 0.001	< 0.001	0.59	<0.001	< 0.001
nPP relative transfer												
Bone mineral content gain	2.60	2.65	2.57	2.61	2.67	2.65	0.15	0.83	0.65	0.76	0.75	0.76
(g/g)												
Tibia ash weight (g/g)	0.337	0.341	0.339	0.362	0.319	0.335	0.012	0.83	0.99	0.07	0.88	0.12
Toe ash weight (g/g)	0.038	0.040	0.042	0.042	0.039	0.040	0.001	0.01	0.03	0.24	0.09	0.54

C = standard commercial levels of Ca and nPP (positive control); M = mid-levels of Ca and nPP; L = low levels; Lin = linear effect; Qua = quadratic effect; ADG = average daily gain; ADFI = average daily feed intake; FCR = feed conversion ratio.

¹For growth performance, n = 10 pens. For bone mineralisation, n = 10 birds.

²First letter corresponds to days 11 to 21. Second letter corresponds to days 22 to 37. See Table 1.

was similar for all treatments. The LC sequence yielded superior bone mineral content as well as tibia and toe ash weights compared with the LM sequence but did not reach the levels obtained with the CC sequence (Lin, P < 0.05). The LM sequence led to poor bone mineral status even though the Ca relative transfer to whole-body, tibia and toe were higher than for CC or LC (Lin, P < 0.001, qua, P < 0.05). Diet had no effect on nPP relative transfer.

Overall performance

The reduced mineral intake had no effect on growth performance from days 0 to 37 (CC *v*. MM *v*. LL, Table 4). In comparison with CC or MM, the LL sequence decreased bone mineral content gain (Lin, Qua, P < 0.001). However, Ca relative transfer to whole-body, tibia and toe were higher in chickens fed LL or MM (Lin, Qua, P < 0.001). These two diets also improved nPP relative transfer to toe (Lin, P < 0.01, Qua, P < 0.01). The duration of the mineral depletion treatment had no effect on growth performance, bone mineral content gain or Ca and nPP relative transfer. The level of Ca and nPP repletion (CC v. LC v. LM contrast) had no effect on growth performance. The degree of repletion that followed the grower phase depletion treatment (i.e. M diet v. C diet during the finisher phase) decreased the bone mineral content gain but also increased the Ca relative transfer to whole-body, tibia and toe (Lin, P < 0.01) and the nPP relative transfer to whole-body (Lin, P < 0.05). The LC and LM treatments both resulted in higher nPP relative transfer to toe than did the control (CC) treatment (Lin, P = 0.001, Qua, P < 0.05).

Discussion

Effect of dietary calcium and non-phytate phosphorus depletion

Reducing dietary nPP and Ca levels during the grower phase did not appear to affect growth performance. In a meta-analysis, Faridi *et al.* (2015) found that 0.60% Ca and 0.30% nPP allow maximal growth performance. At day 21, a decrease of Ca and nPP dietary content led to a decrease of

Valable, Narcy, Duclos, Pomar, Page, Nasir, Magnin and Létourneau-Montminy

								P value ¹					
									Depletio	n	Repletion		
		Diet ²						CC v. MM v. LL		ML v. LL	CC v. LC v. LM		
Parameters	СС	MM	LL	ML	LC	LM	SEM	Lin	Qua	Lin	Lin	Qua	
Growth performance													
ADG (g/d)	64.46	63.97	63.85	63.83	64.78	65.21	0.77	0.42	0.58	0.87	0.34	0.93	
ADFI (g/d)	101.6	102.5	101.7	102.1	102.1	102.5	1.2	0.94	0.73	0.76	0.46	0.95	
FCR	1.570	1.603	1.592	1.599	1.576	1.573	0.018	0.19	0.69	0.81	0.84	0.77	
Mortality (%)	1.49	2.05	2.35	2.26	1.67	1.88	0.91	0.33	0.45	0.82	0.67	0.99	
Bone mineralisation (10 to 37 days)													
Bone mineral content gain (10 to	34.7	32.3	29.2	30.6	34.1	31.3	1.2	< 0.001	< 0.001	0.12	0.01	0.30	
37 days, g/d)													
Ca relative transfer													
Bone mineral content gain (g/g)	0.884	1.125	1.129	1.108	0.999	1.174	0.038	< 0.001	< 0.001	0.65	< 0.0001	0.37	
Tibia ash weight (g/g)	0.072	0.088	0.089	0.090	0.079	0.093	0.003	< 0.001	< 0.001	0.47	< 0.0001	0.16	
Toe ash weight (g/g)	0.008	0.011	0.011	0.011	0.010	0.011	0.001	< 0.001	< 0.001	0.99	< 0.0001	0.55	
nPP relative transfer													
Bone mineral content gain (g/g)	2.183	2.256	2.288	2.244	2.368	2.355	0.082	0.18	0.30	0.88	0.04	0.15	
Tibia ash weight (g/g)	0.177	0.177	0.181	0.183	0.186	0.186	0.007	0.51	0.44	0.32	0.14	0.42	
Toe ash weight (g/g)	0.020	0.021	0.022	0.021	0.023	0.023	0.001	0.002	0.003	0.75	0.001	0.04	

 Table 4 Effect of calcium (Ca) and non-phytate phosphorus (nPP) on broiler chicken overall growth performance and bone mineralisation (days 0 to 37)

C = standard commercial levels of Ca and nPP (positive control); M = mid-levels of Ca and nPP; L = low levels; Lin = linear effect; Qua = quadratic effect; ADG = average daily gain; ADFI = average daily feed intake; FCR = feed conversion ratio.

¹For growth performance, n = 10 pens. For bone mineralisation, n = 10 birds.

²First letter corresponds to days 11 to 21. Second letter corresponds to days 22 to 37. See Table 1.

the body lean content and an increase of body fat content. Phosphorus ingested is fixed at 40% in the soft tissue (Narcy et al., 2015) and will be fixed in priority in the soft tissue over in the bones (Létourneau-Montminy et al., 2015). As a result, a decrease in Ca and nPP content led to a decrease in soft tissue content in the body. As for the body fat content, in the digestive tract, Ca can precipitate with fatty acids, making the subsequent availability of both uncertain. Dietary Ca in excess might reduce carcass fat content by forming such precipitates and thereby interfering with fat emulsification, absorption and deposition (Shafey, 1998). However, body lean and fat content results must be interpreted with caution as the accuracy of DXA analysis for this parameter is lower than that of bone ash determination. Some of the variability could be due to factors that arise during data acquisition, such as the scan programme, mode and bird BW (Swennen et al., 2004).

Between 11 to 21 days, a broiler feed containing 33% less Ca than the NRC-recommended level and 23% less nPP decreased bone mineralisation as shown previously (Rousseau *et al.*, 2013; Hamdi *et al.*, 2015). Faridi *et al.* (2015) showed that in the absence of phytase or vitamin D₃, dietary nPP should be above 3.5 g/kg during the first 21 days in order to maximise tibia ash content. As dietary depletion increased, Ca and nPP relative transfer increased suggesting an increase in apparent digestibility of Ca and phosphorus in the intestine as reported by Rousseau *et al.* (2016) in chicks fed low-phosphorus or Ca diets for 10 days. Studies have

1560

suggested that depletion appears to stimulate the expression of mRNA encoding Ca transporters CALB1, SLC8A1 and ATP2B1 and phosphorus transporters SLC20A1 and SLC34A2 (Centeno *et al.*, 2004; Bar, 2009; Proszkowiec-Weglarz and Angel, 2013) in the intestine and the kidney. Given this adaptation process, more efficient utilisation of both Ca and phosphorus in the intestine may be expected in response to reduced dietary content. Furthermore, the adaptation to low Ca and nPP diet could have occurred primarily in the kidney with an increase in the reabsorption of the minerals. Studies have shown that low plasma Ca and phosphorus concentrations to the stimulation of Ca and phosphorus transporters in the kidney (Proszkowiec-Weglarz and Angel, 2013).

The absence of growth performance alteration when reducing dietary Ca and nPP during days 22 to 37 is in agreement with the results of Wilkinson *et al.* (2014), Delezie *et al.* (2015) and Rousseau *et al.* (2016) for BW gain and feed conversion ratio. A further reduction of Ca and nPP during the finisher phase increased the deficit in bone mineralisation, from -11% on day 21 (C v. L) to -20% on day 37 (CC v. LL) based on tibia ash content. The effect of a decrease in Ca and nPP dietary content on bone mineralisation but not on growth performance further confirmed that the requirements for maximising skeletal development are greater than those for lean mass development (Larbier and Leclercq, 1992). Although it is well known that about 85% of phosphorus is found into bone, in modern broilers that amount is

higher probably due to higher body protein as more than 60% of the phosphorus ingested is deposited in bone with Ca (Narcy *et al.*, 2015). The duration of the depletion treatment (ML *v*. LL) did not impact bone mineralisation (ML and LL led to the same final Ca and nPP relative transfer to whole-body, tibia and toe). We can conclude that the decreased in nPP and Ca content during finisher phase was more crucial to bone development than any adaptation occurring during the decreased in grower phase.

Although the depletion treatment decreased the bone mineralisation values measured on day 37, Ca relative transfer to whole-body, tibia and toe was higher compared with the control treatment. Yan et al. (2005) also demonstrated 20% increases in total Ca apparent absorption measured on day 32 in birds fed such diets during both phases compared with those fed a control diet. The more efficient utilisation of Ca and phosphorus confirms adaptation in depleted animals as shown previously by Rousseau et al. (2016). The absence of significant effects of a decreased in Ca and nPP dietary content on nPP relative transfer to whole-body, tibia, except based on toe ash during the finisher period suggests that Ca may have been the limiting nutrient in our experiment. The relative transfer to whole-body, tibia and toe were similar for ML and LL. Nonetheless, Yan et al. (2005) showed that birds that had been depleted from days 0 to 18 responded better to continued depletion from days 19 to 32 with enhanced Ca and nPP relative transfer to tibia compared with birds receiving a control diet from days 0 to 18 then depleted from days 19 to 32. In our study, the adaptation process may have occurred early in the finisher phase, resulting in complete recovery in bone mineralisation, with a subsequent decline in utilisation efficiency, possibly indicating a transient response.

Effect of dietary calcium and non-phytate phosphorus repletion

The bone mineralisation deficit showed the importance of the repletion phase for restoring the bone mineral criteria to acceptable values (Ashwell and Angel, 2010). The level of repletion (CC v. LC v. LM) did affect bone mineralisation. Despite the lesser degree of mineralisation at the end of the depletion period, increasing the dietary supply of the Ca and nPP to level C or even level M resulted in the same breaking strength on day 37 compared with CC birds. For the other bone mineral status criteria, the LC sequence gave a better result than the LM but LC birds were not able to fully compensate to the levels of the control birds. Rousseau et al. (2013) found that the effects of mineral depletion (0.60% Ca and 0.30% nPP) from days 10 to 21 were erased over days 22 to 35 by switching to a diet containing only slightly more of both minerals (0.70% Ca and 0.35% nPP). This difference could be due to higher Ca:nPP ratio for the control diet during finisher phase in our experiment (2.45) compared with Rousseau et al. (2013, 2.00). Despite the reduced bone mineralisation, Ca relative transfer to whole-body during the finisher period was greater following the LC or LM treatment compared with the control treatment. In contrast, nPP

relative transfer to whole-body, tibia and toe was not affected in our experiment. This confirmed that Ca was the limiting element. Létourneau-Montminy et al. (2008) have reported previously that bone mineralisation in chicks fed a marginally deficient diet for 10 days was restored by replenishing the mineral depletion over the next 11 days as a result of improved bone mineral deposition. Ashwell and Angel (2010) have suggested that an early deficiency could lead to an epigenetic imprinting effect that enhances Ca absorption capability. However, the deficiency began early in the Ashwell and Angel study (90 h after hatching) and even earlier (day 1, lasting through day 18) in a study by Yan et al. (2005), compared with day 11 in our study. A deficiency beginning immediately after hatching might indeed have a genetic and an epigenetic effect. In the absence of such an effect, the gain in Ca and nPP utilisation efficiency might decline over a finisher phase lasting 16 days, as in our experiment. As the LC treatment resulted in the same bone mineral status as the CC treatment did, Ca and nPP utilisation efficiency may have been the same overall in both groups over the entire finisher phase. As LC birds reached the same bone mineral status compared with CC birds, they shared the same efficiency to use Ca and nPP for the remaining of the finisher period. As a result, we might not see a statistical difference over the 16-day period. Based on tibia ash and toe ash, Ca utilisation was more efficient in the LM treatment group than in the LC or CC groups. The LC and LM treatments thus suggest strategies for maximising bone mineralisation. If ~75% of total phosphorus intake occurs during the finisher phase and phosphorus excretion is high at the same time, as reported previously (Rousseau et al., 2013), increasing nPP in the finisher diet appears contrary to our objective of reducing phosphorus excretion. The LM diet seems more appropriate. The depletion and repletion strategy could be improved by including a second depletion towards the end of the finisher phase, considering that absorption will be more efficient due to the first depletion, assuming that the metabolic adaptation is not transient.

Overall, reducing dietary nPP and Ca did not improve the litter score (results not shown). Rousseau *et al.* (2016) had found that reducing Ca content from 9.0 to 7.0 g/kg in the diet improved litter dry matter during a finisher phase with a nPP level of 3.5 g/kg. The acceptable litter quality observed according to Welfare Quality (2009) for all of the diets in our experiment could explain the lack of effect (Tuyttens *et al.*, 2015).

Assessment of bone mineralisation

Assessing bone mineral status is critical to the study of the mineral adequacy of diets. Bone physical and mechanical properties such as strength and the structure of bone tissue reflect overall skeletal health (Bradbury *et al.*, 2014). We used several criteria to evaluate the impact of dietary strategies on bone mineralisation. The bone mineral content assessed by DXA scanning demonstrated to discriminate the treatments as well as the tibia and toe ash measurements. Whole-carcass bone mineral content was correlated best with tibia ash weight, followed by the toe ash mass (results not

Valable, Narcy, Duclos, Pomar, Page, Nasir, Magnin and Létourneau-Montminy

shown). These results assessed the validity of DXA scanning in broilers experiment. This technology had been applied previously to broilers by only few authors (Angel et al., 2006; Shahnazari et al., 2007). It present the advantage, compared with traditional measurements, to provide a rapid and noninvasive measurement of whole-body bone mineralisation. We also assess the advantage to use toe instead of tibia for ash measurements. In our experiment, tibia ash and toe ash were poorly correlated, making it difficult to consider toe ash as a good proxy measurement of bone mineralisation. Shastak et al. (2012) found toe ash to be highly correlated with tibia ash in 3-week-old broilers ($R^2 = 0.94$). However, these authors used all toes for the measurement rather than only the middle toe as we did. They also found a stronger correlation between tibia ash and whole foot ash compared with toe ash. The increased accuracy in other experiments for toe ash could be due to the greater number of bones.

Although breaking strength responded to the dietary treatments the same way as other bone mineralisation criteria did (bone mineral content, tibia and toe ash weights), it showed higher variability (coefficients of variation of 6.55% v. 3.39%, 3.55% and 3.32%, respectively). Ravindran *et al.* (1995) did not consider tibia breaking strength as a good indicator of phosphorus bioavailability, as it appeared to be more sensitive than other bone mineral criteria to variations in the measurement technique (handling, shear force, crosshead speed).

In conclusion, this study confirms that it is possible to reduce dietary Ca and nPP levels below the NRC (1994) recommendations during the grower and finisher phases without affecting growth performance. Despite decreased bone mineralisation when mineral depletion occurs during the grower phase, the deficit is restored if the minerals are replenished with control diet during the finisher phase. However, the dietary Ca and nPP levels in the repletion diet determine the final levels of bone mineralisation, the control (commercial conditions) level resulting in better mineralisation than an intermediate level. Bone mineralisation under these conditions benefits from Ca and nPP relative transfer, particularly Ca relative transfer in the finisher phase, being increased in response to grower phase conditions.

Acknowledgements

The authors gratefully acknowledge Nutreco Canada Inc. for its financial support and for conducting the experiment. The authors are grateful to the Sherbrooke centre of Agriculture and Agri-food Canada, Marcel Marcoux and Nancy Bolduc for their technical support in performing the laboratory analysis.

References

Angel R, Saylor WW, Mitchell A, Powers W, Applegate TJ and Dhandu AS 2006. Effect of dietary phosphorus, phytase and 25-hydroxycholecalciferol on broiler chicken bone mineralisation, litter phosphorus, and processing yields. Poultry Science 85, 1200–1211.

Ashwell CM and Angel R 2010. Nutritional genomics: a practical approach by early life conditioning with dietary phosphorus. Revista Brasileira de Zootecnia 39, 268–278.

Applegate TD and Angel R 2008. Phosphorus requirements for poultry. Fact sheet for natural resources conservation service feed management 592 Practice Standard, Education Project, USDA CIG project, West Lafayette, IN, USA.

Bar A 2009. Calcium transport in strongly calcifying laying birds: mechanisms and regulation. Comparative Biochemistry and Physiology (Part A) 152, 447–469.

Bradbury EJ, Wilkinson SJ, Cronin GM, Thomson PC, Bedford MR and Cowieson AJ 2014. Nutritional geometry of calcium and phosphorus nutrition in broiler chicks. Growth performance, skeletal health and intake arrays. Animal 8, 1071–1079.

Canadian Council on Animal Care 2009. Guidelines on the care and use of farm animals in research, teaching and testing. CCAC, Ottawa, ON, Canada.

Centeno VA, Diaz de Barboza GA, Marchionattia AN, Alisioa AE, Dallorsob ME, Nasif R and Tolosa de Talamonia NG 2004. Dietary calcium deficiency increases Ca²⁺ uptake and Ca²⁺ extrusion mechanisms in chick enterocytes. Comparative Biochemistry and Physiology 139, 133–141.

Delezie E, Bierman K, Nollet L and Maertens L 2015. Impacts of calcium and phosphorus concentration, their ratio, and phytase supplementation level on growth performance, foot pad lesions, and hock burn of broiler chickens. Journal of Applied Poultry Research 24, 115–126.

Faridi A, Gitoee A and France J 2015. A meta-analysis of the effects of nonphytate phosphorus on broiler performance and tibia ash concentration. Poultry Science 94, 2753–2762.

Hamdi M, López-Vergé S, Manzanilla EG, Barroeta AC and Pérez JF 2015. Effect of different levels of calcium and phosphorus and their interaction on the performance of young broilers. Poultry Science 94, 2144–2151.

Larbier M and Leclercq B 1992. Nutrition et alimentation des volailles. Institut National de la Recherche Agronomique Editions, Paris, France.

Létourneau-Montminy MP, Lescoat P, Narcy A, Sauvant D, Bernier JF, Magnin M, Pomar C, Nys Y and Jondreville C 2008. Effects of reduced dietary calcium and phytase supplementation on calcium and phosphorus utilisation in broilers with modified mineral status. British Poultry Science 49, 705–715.

Létourneau-Montminy MP, Narcy A, Dourmad JY, Crenshaw TD and Pomar C 2015. Modeling the metabolic fate of dietary phosphorus and calcium and the dynamics of body ash content in growing pigs. Journal of Animal Science 93, 1200–1217.

Létourneau-Montminy MP, Narcy A, Lescoat P, Bernier JF, Magnin M, Pomar C, Nys Y, Sauvant D and Jondreville C 2010. Meta-analysis of phosphorus utilisation by broilers receiving corn-soyabean meal diets: influence of dietary calcium and microbial phytase. Animal 4, 1844–1853.

Mitchell AD, Rosebrough RW and Conway JM 1997. Body composition analysis of chickens by dual energy X-ray absorptiometry. Poultry Science 76, 1746–1752.

Narcy A, Rousseau X, Même N, Magnin M and Nys Y 2015. Impact of dietary phosphorus and calcium levels on their deposition in soft and bone tissues in chickens. In Proceedings of the 20th European Symposium of Poultry Nutrition, 24–27 August 2015, Prague, CZR, p. 217.

National Research Council (NRC) 1994. Nutrient requirements of poultry, 9th revised edition. National Academy Press, Washington, DC, USA.

Proszkowiec-Weglarz M and Angel R 2013. Calcium and phosphorus metabolism in broilers: effect of homeostatic mechanism on calcium and phosphorus digestibility. Journal of Applied Poultry Research 22, 609–627.

Ravindran V, Kornegay ET, Potter LM, Ogunabameru BO, Welten MK, Wilson JH and Potchanakorn M 1995. An evaluation of various response criteria in assessing biological availability of phosphorus for broilers. Poultry Science 74, 1820–1830.

Rousseau X, Même N, Magnin M, Couty M, Bordeau T and Narcy A 2013. Vers une utilisation efficace des ressources minérales. Paper presented at the 10th Journées de la Recherche Avicole et Palmipèdes à Foie Gras, 26–28 March 2013, La Rochelle, France, pp. 650–654.

Rousseau X, Valable AS, Létourneau-Montminy MP, Même N, Godet E, Magnin M, Nys Y, Duclos MJ and Narcy A 2016. Adaptive response of broilers to dietary phosphorus and calcium restrictions. Poultry Science 95, 2849–2860.

Schreiweis MA, Orban JI, Ledur MC, Moody DE and Hester PY 2005. Validation of dual-energy X-ray absorptiometry in live white leghorns. Poultry Science 84, 91–99.

Shafey TM 1998. Effects of dietary calcium, phosphorus, biotin and fat on the performance and nutrient utilisation of meat chickens. Agriculture Science 10, 121–132.

Shahnazari M, Lang DH, Fosmire GJ, Sharkey NA, Mitchell AD and Leach RM 2007. Strontium administration in young chickens improves bone volume and architecture but does not enhance bone structural and material strength. Calcified Tissue International 80, 160–166.

Shastak Y, Witzig M, Hartung K, Bessei W and Rodehutscord M 2012. Comparison and evaluation of bone measurements for the assessment of mineral phosphorus sources in broilers. Poultry Science 91, 2210–2220.

Statistical Analysis System Institute Inc 2003. SAS 9.0 user's guide. SAS Institute Inc, Cary, NC, USA.

Swennen Q, Janssens GP, Geers R, Decuypere E and Buyse J 2004. Validation of dual-energy x-ray absorptiometry for determining *in vivo* body composition of chickens. Poultry Science 83, 1348–1357.

Tuyttens FA, Federici JF, Vanderhasselt RF, Goethals K, Duchateau L, Sans ECO and Molento CFM 2015. Assessment of welfare of Brazilian and Belgian broiler flocks using the Welfare Quality protocol. Poultry Science 94, 1758–1766.

Welfare Quality 2009. Welfare quality assessment protocols for poultry (broilers, laying hens). Welfare Quality Consortium, Lelystad, the Netherlands.

Wilkinson SJ, Bradbury EJ, Bedford MR and Cowieson AJ 2014. Effect of dietary non-phytate phosphorus and calcium concentration on calcium appetite of broiler chicks. Poultry Science 93, 1695–1703.

Yan F, Angel R, Ashwell C, Mitchell A and Christman M 2005. Evaluation of the broiler's ability to adapt to an early moderate deficiency of phosphorus and calcium. Poultry Science 84, 1232–1241.